



Elfabrio Home Infusion Nursing Order Form



Please fax completed form to Chiesi Total CareSM staff at 1-636-355-3610.

PATIENT INFORMATION

Patient Name (Last, First) _____
 Social Security # _____ - _____ - _____ Sex: Male Female Date of Birth ____/____/____ (mm/dd/yyyy)
 Address _____ City _____ State _____ ZIP _____
 Primary Phone (Required) _____ Cell Phone _____ Language: English Other _____

MEDICAL INFORMATION

Diagnosis: Fabry (-Anderson) disease E75.21
 Height _____ inches or _____ cm Weight _____ lb or _____ kg Allergies: None Specify _____
 Prior Therapies _____
 Current Medications _____
 History of Infusion-Related Reactions _____

ELFABRIO® (PEGUNIGALSIDASE ALFA-IWXJ) 20 mg/10 mL PRESCRIPTION

Dosage — Elfabrio (pegunigalsidase alfa-iwxj) 20 mg/10 mL vial
 Total Dose (mg*) _____ Route of Administration IV _____ Frequency* _____
 Number of Refills† _____ Infusion Rate (mL/hour) _____
 Please list any additional treatment information, including follow-up evaluations: _____

*The recommended dosage is 1 mg/kg of body weight every 2 weeks, administered as an intravenous infusion.
 †Quantity sufficient for a 28-day supply.

NURSING ORDERS

- The recommended dose is 1 mg/kg of body weight
 - Number of vials per infusion (round up to a whole # if the calculated # of vials is a fraction) = Patient dose (mg) ÷ 20 mg (contents of 1 vial)
 - Prior to adding the volume of Elfabrio required for the dose, remove the equal volume of 0.9% Sodium Chloride Injection from the infusion bag
 - Infusion duration is calculated individually, with a minimum infusion duration of 90 minutes
- Monitor vital signs at the start of the infusion, at the end of the infusion and observation period, and PRN _____
- Intravenous access and flush orders:
 - Peripheral IV line:
 - Before Infusion: 0.9% Sodium Chloride Injection 3-5 mL
 - After Infusion: 0.9% Sodium Chloride Injection 3-5 mL
 - Implanted port/Central line:
 - Before Infusion: 0.9% Sodium Chloride Injection 5-10 mL
 - After Infusion: 0.9% Sodium Chloride Injection 5-10 mL and Heparin (100 units/mL) 5 mL

Patient's Actual Weight	Total Infusion Volume
up to 70 kg	<input type="checkbox"/> 150 mL 0.9% NaCl
70 - 100 kg	<input type="checkbox"/> 250 mL 0.9% NaCl
>100 kg	<input type="checkbox"/> 500 mL 0.9% NaCl

• Infusion rate may be increased if the patient tolerates the initial 4-6 Elfabrio infusions. Infusion rate may be slowed in case of hypersensitivity reaction or an infusion-associated reaction

If a severe reaction occurs, immediately discontinue Elfabrio, initiate appropriate medical care, and contact the prescriber.
If a mild to moderate reaction occurs, consider slowing or temporarily withholding Elfabrio, initiate appropriate medical care, and contact the prescriber.

PHYSICIAN/OFFICE INFORMATION

Prescriber's Name (Print) _____ Practice/Group Name _____
 Address _____ Suite _____
 City _____ State _____ ZIP _____
 Office Contact Person _____ Office Phone _____
 Emergency Cell Phone _____ Office Fax _____
 License # _____ NPI # _____

Please see Important Safety Information for Elfabrio, including Boxed Warning concerning hypersensitivity reactions including anaphylaxis, on page 3 of this form and accompanying Full Prescribing Information.

Please call Chiesi Total CareSM at 1-833-656-1056 if you have any questions regarding this form.

MEDICATION ORDERS

Select premedication(s) needed for this administration:

Oral (PO) medications to be obtained and self-administered by patient

- EMLA™ Cream
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Methylprednisolone
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Acetaminophen
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Famotidine
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Diphenhydramine (50 mg/mL)
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Other: _____
Dose _____ Route _____
Physician Directions _____
Quantity _____ Refills _____

Additional orders:

Select PRN medications:

Oral (PO) medications to be obtained and self-administered by patient

- Methylprednisolone
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Acetaminophen
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Famotidine
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Diphenhydramine
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Albuterol sulfate inhalation aerosol for oral inhalation
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Epinephrine auto-injector
Dose _____
Physician Directions _____
Quantity _____ Refills _____
- Other: _____
Dose _____ Route _____
Physician Directions _____
Quantity _____ Refills _____
- Emergency kit:
 - Epinephrine auto-injector
 - PPE for administering CPR

MANDATORY OFFICE CHECKLIST

 Please confirm that you have completed each of the following steps:

- The patient has had successful infusions in an outpatient setting
- The patient is medically stable and safe for home infusion therapy
- The patient has been prescribed an epinephrine auto-injector
- The patient lives in an area where emergency medical services are available

By signing below, I certify that I am the prescribing provider mentioned above, that I am part of the Chiesi Total CareSM Program, that the therapy described above is medically necessary, and that all the medical necessity information is true, accurate, and complete. The patient's records contain supporting documentation that substantiates the utilization and medical necessity of the products marked above. I provide permission to use my personal information and the personal information of the patient provided above to facilitate this request and complete any regulatory or legal requirements associated with this request. I understand that the personal information provided herein may be shared with Chiesi, successors, and their agents and service providers as needed to support this request. I also attest that I have obtained the patient's authorization to release the above information and such other personal information as may be necessary for the Chiesi Total Care Program and/or their agents and service providers. If the patient is 18 years old or younger, I attest that I have obtained permission from the patient's legal guardian. If my patient is eligible for free product, I understand that receiving free product is not contingent on any purchase obligations. I also understand that no free product may be submitted for reimbursement to any payer, including Medicare and Medicaid; nor may I bill any payer for administration of such product. I understand that any falsification, omission, or concealment of material fact may result in criminal liability.

Treating Physician's Signature

Date

Please see Important Safety Information for Elfabrio, including Boxed Warning concerning hypersensitivity reactions including anaphylaxis, on page 3 of this form and accompanying Full Prescribing Information.

Please call Chiesi Total CareSM at 1-833-656-1056 if you have any questions regarding this form.

Indication

Elfabrio[®] (pegunigalsidase alfa-iwxj) is indicated for the treatment of adults with confirmed Fabry disease.

Important Safety Information

WARNING: HYPERSENSITIVITY REACTIONS INCLUDING ANAPHYLAXIS

Patients treated with Elfabrio have experienced hypersensitivity reactions, including anaphylaxis. Appropriate medical support measures, including cardiopulmonary resuscitation equipment, should be readily available during Elfabrio administration. If a severe hypersensitivity reaction (eg, anaphylaxis) occurs, discontinue Elfabrio immediately and initiate appropriate medical treatment. In patients with severe hypersensitivity reaction, a desensitization procedure to Elfabrio may be considered.

Prior to Elfabrio administration, consider pretreating with antihistamines, antipyretics, and/or corticosteroids. Inform patients and caregivers of the signs and symptoms of hypersensitivity reactions and infusion-associated reactions (IARs), and instruct them to seek medical care immediately if such symptoms occur.

- If a severe hypersensitivity reaction (including anaphylaxis) or severe IAR occurs, immediately discontinue Elfabrio administration and initiate appropriate medical treatment.
- If a mild to moderate hypersensitivity reaction or IAR occurs, consider slowing the infusion rate or temporarily withholding the dose.

In clinical trials, 20 (14%) Elfabrio-treated patients experienced hypersensitivity reactions. Four Elfabrio-treated patients (3%) experienced anaphylaxis reactions that occurred within 5 to 40 minutes of the start of the initial infusion. The signs and symptoms of hypersensitivity reactions and anaphylaxis included headache, nausea, vomiting, throat tightness, facial and oral edema, truncal rash, tachycardia, hypotension, rigors, urticaria, intense pruritus, moderate upper airway obstructions, macroglossia, and mild lip edema.

In clinical trials, 41 (29%) Elfabrio-treated patients experienced one or more infusion-associated reactions, including hypersensitivity, nausea, chills, pruritus, rash, chest pain, dizziness, vomiting, asthenia, pain, sneezing, dyspnea, nasal congestion, throat irritation, abdominal pain, erythema, diarrhea, burning sensation, neuralgia, headache, paresthesia, tremor, agitation, increased body temperature, flushing, bradycardia, myalgia, hypertension, and hypotension.

A case of membranoproliferative glomerulonephritis with immune depositions in the kidney was reported during clinical trials. Monitor serum creatinine and urinary protein-to-creatinine ratio. If glomerulonephritis is suspected, discontinue treatment until a diagnostic evaluation can be conducted.

When switching to Elfabrio from a prior enzyme replacement therapy, the risk of hypersensitivity reactions and infusion-associated reactions may be increased in certain patients with pre-existing anti-drug antibodies (ADAs). Consider monitoring IgG and IgE ADAs and clinical or pharmacodynamic response (eg, plasma lyso-Gb3 levels).

The most common adverse reactions ($\geq 15\%$) were infusion-associated reactions, nasopharyngitis, headache, diarrhea, fatigue, nausea, back pain, pain in extremity, and sinusitis.

Please see accompanying [Full Prescribing Information](#) for Elfabrio.