



For adult and pediatric patients 6 months of age and older with dystrophic or junctional epidermolysis bullosa (EB)

Getting Started With FILSUVEZ

This guide will help you better understand your treatment.

Important Safety Information

What is FILSUVEZ (birch triterpenes) topical gel for?

FILSUVEZ is a prescription medicine used on the skin to treat wounds that may happen with dystrophic and junctional epidermolysis bullosa (EB) in adults and children 6 months of age and older.

It is not known if FILSUVEZ is safe and effective in children younger than 6 months of age.

Please see [Important Safety Information](#) throughout. Please see full [Product Information](#) and [Patient Information](#).



Filsuvez[®]
(birch triterpenes) topical gel

What is FILSUVEZ?




FILSUVEZ is a prescription medicine used on the skin to support wound healing in adults and children with dystrophic epidermolysis bullosa (DEB) or junctional epidermolysis bullosa (JEB).

What should I expect from treatment with FILSUVEZ?

You should experience wound healing and closure in about 45 days, as seen in a phase 3 study of FILSUVEZ conducted in 223 EB patients—195 with DEB and 26 with JEB.

45 days

In that same study, 41.3% of people using FILSUVEZ achieved wound closure in 45 days compared with 28.9% of those using the placebo gel.



Important Safety Information (cont'd)

What is the most important safety information I should know?

FILSUVEZ may cause a serious side effect of allergic reactions.

Allergic reactions and skin reactions to FILSUVEZ may include the following symptoms: red itchy bumps (hives), skin rash, redness or itching. If you get any of these symptoms, stop using FILSUVEZ right away and call your healthcare provider.

Please see [Important Safety Information](#) throughout.

Please see full [Product Information](#) and [Patient Information](#).

Filsuvez[®]
(birch triterpenes) topical gel

How should I apply FILSUVEZ?



Step 1



Clean all wounds and apply a generous layer of FILSUVEZ to each nonadhesive wound dressing or directly to the wounds with a clean or gloved hand. **Do not** rub in the gel.

Step 2



Spread FILSUVEZ, ensuring that each layer of gel is about the thickness of a credit card.

Step 3



Apply dressings to all wounds and reapply FILSUVEZ with each dressing change until all wounds are fully healed.

Always use FILSUVEZ exactly as instructed by your healthcare provider.



Watch a demonstration on how FILSUVEZ is applied.

Simply scan the QR code or visit www.filsuvez.com to learn more.

Important Safety Information (cont'd)

What are the possible side effects of FILSUVEZ?

The most common side effect of FILSUVEZ is application site reactions, such as pain and itchy skin.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088

Please see [Important Safety Information](#) throughout. Please see full [Product Information](#) and [Patient Information](#).

Filsuvez[®]
(birch triterpenes) topical gel

What else should I know about using FILSUVEZ?



FILSUVEZ should be applied evenly to **all wounds** at **every wound dressing change**. You may use the same tube on **multiple wounds** during a single dressing change.



One tube of FILSUVEZ covers about two 5x4-inch bandages, nearly the same area as a business-sized envelope. Use the Coverage Calculator found in your Starter Kit to determine how many tubes you need for your wounds at each dressing change, but **do not place this tool directly on your wounds**.



Each tube of FILSUVEZ should only be used once. Once the tube has been opened, apply the gel to all of your wounds immediately and then dispose of any remaining gel and the tube in your household trash.



If the tube runs out during a dressing change, open a new tube and continue applying the gel until all of your wounds are covered.



FILSUVEZ should be stored at 68°F to 77°F (20°C to 25°C). Do not freeze.



Do not use FILSUVEZ in or around the eyes or mucous membranes (mouth, vagina, or anus). If accidental contact does occur, immediately wash with clean water and contact your doctor if you have any discomfort.

Important Safety Information (cont'd)

How should I take FILSUVEZ?

See the detailed “Instructions for Use” that comes with FILSUVEZ for information on how to apply FILSUVEZ.

Use FILSUVEZ exactly as your healthcare provider tells you to use it.

Please see [Important Safety Information](#) throughout. Please see full [Product Information](#) and [Patient Information](#).

Filsuvez[®]
(birch triterpenes) topical gel

What are the possible side effects of FILSUVEZ?



FILSUVEZ may cause serious side effects

If you get an allergic reaction, stop using FILSUVEZ immediately and contact your doctor.

- The most common side effect of FILSUVEZ is application site reactions, such as pain and itchy skin
- If the wounds you are treating with FILSUVEZ become infected, **stop treatment** and contact your doctor
 - Possible signs of infection include, but are not limited to, the wound becoming red, warm, swollen, painful, and/or draining yellow or greenish fluids (pus)

These are not all the possible side effects of FILSUVEZ.

Call your doctor for medical advice about side effects.
You may report side effects to the FDA at 1-800-FDA-1088
(1-800-332-1088).

Important Safety Information (cont'd)

How should I take FILSUVEZ? (cont'd)

The tube of FILSUVEZ is for one-time use only. After the tube has been opened, apply the gel right away. Throw away any remaining gel and the tube after use.

Please see [Important Safety Information](#) throughout.
Please see full [Product Information](#) and [Patient Information](#).

**Filsuvez**[®]
(birch triterpenes) topical gel

What support is available with FILSUVEZ?



One-Stop Patient Support

Chiesi Total CareSM is a personalized and comprehensive support program designed to help patients using FILSUVEZ and their healthcare providers.



The Chiesi Total Care Program Assists With:



Insurance Consultation and Financial Support

- Help with understanding health insurance benefits and coverage for FILSUVEZ
- Information on financial assistance options for those who are eligible



Treatment Support and Logistics Coordination

- Home delivery of your FILSUVEZ prescription each month
- Help staying on track with prescription refill reminders



Nurse Support and Personalized Education

- Customized education on EB and how to use FILSUVEZ available via phone, virtually, or in person with a Registered Nurse



Dedicated Total Care Team

- A helpful, caring team is here to help, with Pharmacists, Patient Service Coordinators, Reimbursement Support Specialists, and Nursing Support

Call **1-833-670-6464**, Mon-Fri 8am to 8pm ET or visit www.chiesitotalcare.com to learn more.

Important Safety Information (cont'd)

How should I take FILSUVEZ? (cont'd)

Do not use around or get FILSUVEZ in the eyes, or mucous membrane areas, examples are mouth, vagina or anus.

Please see **Important Safety Information** throughout.
Please see full **Product Information** and **Patient Information**.

Filsuvez[®]
(birch triterpenes) topical gel



For advocacy information and other related support, please visit:



debra of America, an organization dedicated to helping the EB community since 1980. Learn more at www.debra.org



EB Research Partnership (EBRP), a global organization dedicated to funding research to treat and cure EB. Learn more at www.ebresearch.org

Important Safety Information (cont'd)

How should I take FILSUVEZ? (cont'd)

If you get FILSUVEZ in your eyes or mucous membrane area, rinse with clean water right away. Contact your healthcare provider if you have any discomfort.

Please see [Important Safety Information](#) throughout. Please see full [Product Information](#) and [Patient Information](#).





For adult and pediatric patients 6 months of age and older with dystrophic or junctional epidermolysis bullosa (EB)

Reimagine Living With EB



FILSUVEZ is an FDA-approved topical gel that supports wound healing in people with DEB and JEB.

For more information, visit www.filsuvez.com or simply scan the QR code.

Important Safety Information (cont'd)

How should I take FILSUVEZ? (cont'd)

If the wounds you are treating with FILSUVEZ become infected, **stop treatment** and contact your healthcare provider. Signs or symptoms of infection may include the wound becoming red, warm, swollen, painful or drains yellow or greenish fluid (pus).

Please see **Important Safety Information** throughout.

Please see full **Product Information** and **Patient Information**.



©2024 Chiesi USA. All rights reserved.
Filsuvez® is a registered trademark owned by the Chiesi Group.
Chiesi Total CareSM is a service mark owned by Chiesi Farmaceutici S.p.A.
Other trademarks are the property of their respective owners.
PP-FZ-0088 V2.0

Filsuvez[®]
(birch triterpenes) topical gel