



What is JUXTAPID?

JUXTAPID is a medicine that your doctor gives you to lower cholesterol. It helps those with a rare genetic condition called homozygous familial hypercholesterolemia (HoFH). Doctors use JUXTAPID with a low-fat diet and other cholesterol-lowering treatments.

Due to the risk of liver problems, JUXTAPID is only for people with HoFH. It is not known if JUXTAPID is safe and effective for people with high cholesterol who do not have HoFH such as those with heterozygous familial hypercholesterolemia (HeFH). HeFH type is less severe, and only one parent passes it down.

Risk of liver problems with JUXTAPID

JUXTAPID can lead to liver problems. This includes higher liver enzymes and fat buildup in the liver.

Your doctor will order blood tests to check your liver at regular intervals:

- before you start JUXTAPID
- if they increase your dose
- monthly during the first year
- every 3 months after the first year

Blood test results show if your liver enzyme levels are too high. Enzyme levels higher than normal can be an early sign of liver problems. If your tests show liver problems, your doctor might order more tests. They may also lower your dose or stop it.

JUXTAPID may cause other side effects. Talk to your doctor about the risks of these and any other questions you have about taking JUXTAPID.

What should I do?

Before you start treatment with JUXTAPID, tell your doctor if you have had liver problems, especially with other medications.

If you are taking JUXTAPID, let your caregiver and doctor know right away if any of these happen. They could be signs of liver issues:

- nausea (feeling sick), vomiting (being sick), or stomach pain that gets worse, changes, or does not go away
- fever
- flu-like symptoms
- your eyes or skin become yellow
- feeling more tired than usual

JUXTAPID can cause nausea, vomiting, stomach pain, diarrhea, gas, and burping. This is more likely when you don't follow your low-fat diet. Some side effects may be symptoms of liver problems. If they occur, tell your caregiver or doctor.

You should not use certain other medicines at the same time as JUXTAPID. Tell your caregiver(s) and doctor about all the medicines you take. This includes both prescription and over-the-counter drugs. Some medicines, when taken together, can overwork the liver and cause problems.

You should take your JUXTAPID capsule whole with water at least 2 hours after the evening meal every day. If you cannot swallow the capsule, you can open it, sprinkle its contents on 1 tablespoon of applesauce or mashed banana to take it.

Take your vitamin supplements every day.



Instructions for Prescribers

The form must be signed by both the prescriber and the parent or legal guardian* of any patient under the age of 18 years. Provide a signed copy of the form to the patient.

PARENT OR GUARDIAN AND PATIENT ACKNOWLEDGEMENT

Patient First Name: _____ Middle Initial: ____ Last Name: _____
City: _____ State: _____ Zip: _____
Phone: _____ Email: _____
Date of Birth: _____ Patient Gender: [] Male [] Female [] Decline to Identify

I have received the Patient Guide from my prescriber. I understand that:

- JUXTAPID is used along with a low-fat diet, exercise, and other low density lipoprotein cholesterol (LDL-C) lowering medicines to reduce LDL-C in adults and children 2 years and older with a type of high cholesterol called homozygous familial hypercholesterolemia (HoFH).
JUXTAPID may cause serious side effects including liver problems such as increased liver enzymes or increased fat in the liver.
Because of these side effects, JUXTAPID is only for people with homozygous familial hypercholesterolemia (HoFH).
I will enroll in the JUXTAPID REMS by completing this Patient-Prescriber Acknowledgement Form.

- I will need to have blood tests to check my liver:
- before I start JUXTAPID
- if my dose is increased
- monthly during the first year
- every three months after the first year
If my tests show liver problems, my doctor may lower my dose of JUXTAPID or stop it.
I will tell my doctor if I have any of the following symptoms:
- nausea, vomiting, or stomach pain that gets worse, changes, or does not go away
- fever
- flu-like symptoms
- yellowing of my eyes or skin
- feeling more tired than usual

*Parent or Guardian Signature: _____ Date: _____
Parent/Guardian First Name: _____ Parent/Guardian Last Name: _____

PRESCRIBER ACKNOWLEDGEMENT

Prescriber First Name: _____ Middle Initial: ____ Last Name: _____
Address: _____
City: _____ State: _____ Zip: _____
Office Phone: _____ NPI #: _____

- I have counseled the parent/guardian and patient on the indication and risks of JUXTAPID, including the risk of liver problems, and the need for periodic liver monitoring.
I have reviewed the Patient Guide with the parent/guardian and patient and provided a signed copy of this form to the parent/guardian (and patient when appropriate).
I discussed all concerns and answered all questions the parent/guardian and patient had about treatment with JUXTAPID.

Prescriber Signature: _____ Date: _____

IMPORTANT

REVIEW TO ENSURE ALL FIELDS ARE COMPLETED | FAX TO 1-855-898-2498

If you have any questions, please contact the JUXTAPID REMS Coordinating Center.
Phone: 1-85-JUXTAPID (1-855-898-2743) | Fax: 1-855-898-2498 | www.juxtapidREMSprogram.com

