Juxtapid[®] Chiesi Total Care[®] Patient Support Services Program Terms and Conditions

To enroll in Chiesi Total Care (the "Program") and to assess eligibility for patient support services of the Program, patient must complete the Program Enrollment and Authorization Form and have a valid prescription for an eligible product of the Program. Additional documentation may be required. The patient must be a resident of the United States or one of its territories. If the patient is incapable of acting on their own behalf or if the patient is under 18 years old, enrollment into the Program may be completed by another person acting on their behalf (such as a caregiver).

A patient who receives healthcare benefits under any plan or program funded in whole or in part by federal or state governments including Medicare, Medicaid, TRICARE, Veterans Affairs (VA), State Prescription Assistance Plans (SPAPs) (other than health insurance for federal government employees), or any state healthcare program such as Medicaid, Children's Health Insurance Program, programs funded under Maternal and Child Health Program, or programs funded under Social Services Block Grant (collectively, "Governmentfunded Plans") is not eligible for the financial patient support services of the Program. A patient covered under a commercial health plan purchased through a health insurance marketplace or exchange is not a Government-funded Plan beneficiary even if the costs of such coverage are subsidized by the federal government. If a change in prescription drug coverage should occur, the patient must notify the Program; such change may affect eligibility for the support services provided in the Program. Patients who have been prescribed a product for an indication that is not consistent with the US Food and Drug Administration-approved labeling will not be eligible for financial patient support services offered through the Program.

Patients residing in or receiving treatment in certain states may not be eligible for certain patient support services of the Program. Patients may not seek reimbursement for value received from the Program. The Program does not obligate the use of any specific medication or healthcare provider. Patients who receive treatment or reside in Massachusetts or Rhode Island are not eligible for co-pay assistance for infusion services or routine testing services.

Chiesi Total Care may recommend contacting an independent financial assistance foundation. Independent financial assistance foundations have their own rules for eligibility. Chiesi USA does not fund independent financial assistance foundations, nor does Chiesi Total Care have involvement or influence in independent foundation decision making or eligibility criteria and does not know if a foundation will be able to help you. Chiesi Total Care can only refer you to a foundation that supports your disease state. This information is provided as a resource for you. Chiesi Total Care does not endorse or show preference for any foundation. The foundations recommended to you may not be the only ones that might be able to help you.

Chiesi Patient Education Liaisons ("PELs") may be available to assist you with disease education, provide relative educational or informational resources, and to answer questions you may have about your disease. Chiesi Field Reimbursement Managers ("FRMs") may be available to assist you with your product prescription drug coverage, including prior authorization, appeals, and denials.

PELs and FRMs are employees of Chiesi USA, Inc. PELs and FRMs are not healthcare providers and are not part of your healthcare team. PELs or FRMs will not provide medical care or advice. All treatment decisions should be made by you and your treating healthcare professional. To assist you, PELs and FRMs may need your Information. If you choose to opt out of services by PELs and FRMs, you may do so at any time. Please see Chiesi's Privacy Policy at www.chiesiusa.com/privacy.

Program benefits may not be sold, purchased, traded, or offered for sale, purchase, or trade. The Chiesi Total Care patient support services are not valid where prohibited by law, taxed, or otherwise restricted. Offer subject to change or discontinuance without notice. Restrictions, including monthly maximums, may apply. This is not health insurance.

This is a voluntary program. Patients who choose not to enroll in the Program will be able to receive medication. Patients may participate in Chiesi Total Care without participating in a patient support services program of Chiesi Total Care. After enrolling in Chiesi Total Care, participants may opt out by contacting the Program, as outlined in the Chiesi Total Care Enrollment and Authorization Form. Patients must renew their eligibility by December 31 of each year to continue to receive support under the Program.

By participating in the Program, participants acknowledge that they understand and agree to comply with the Program Terms and Conditions.

Important Safety Information

Indication

Juxtapid is a prescription medicine used along with diet and other lipid lowering treatments, including low-density lipoprotein (LDL) apheresis where available, in adults with homozygous familial hypercholesterolemia (HoFH) to reduce LDL ("bad") cholesterol, total cholesterol, a protein that carries bad cholesterol in the blood (apolipoprotein B), and non-high-density lipoprotein cholesterol (non-HDL-C).

Studies have not been conducted to tell us whether Juxtapid can help prevent problems from high cholesterol, such as heart attack, stroke, death, or other health problems. Studies have also not been conducted to tell us whether Juxtapid is safe for use in people with high cholesterol who do not have HoFH, including those with heterozygous familial hypercholesterolemia (HeFH).

Juxtapid may cause serious side effects, including: Liver Problems

- Juxtapid can cause liver problems such as increased liver enzymes or increased fat in the liver. For this reason, your doctor should do blood tests to check your liver before you start Juxtapid and while you are taking Juxtapid (especially if your dose is increased). If your tests show signs of liver problems, your doctor may lower your dose of Juxtapid or stop it altogether.
- You should tell your doctor if you have had liver problems in the past, including liver problems while taking other medicines.
- Stomach problems can also be a symptom of liver problems. Tell your doctor right away if you have nausea; vomiting or stomach pain that gets worse, does not go away, or changes; fever; yellowing of your eyes or skin; feeling more tired than usual; or having flu-like symptoms while taking Juxtapid because these may be signs of liver problems.
- Do not drink more than 1 alcoholic drink per day while taking Juxtapid.

Harm to your unborn baby:

- Do not take Juxtapid if you are pregnant, think you may be pregnant, or are planning to become pregnant.
- You should have a negative pregnancy test result before you can start on Juxtapid. Use effective birth control while taking Juxtapid. If you become pregnant while taking Juxtapid, stop taking Juxtapid and call your doctor right away.

You should not take Juxtapid if you:

- Are taking medications known as moderate or strong CYP3A4 inhibitors (for example, certain medications used to treat bacterial, fungal, or viral infections, as well as certain medications used to treat depression, high blood pressure, or angina). These medications may affect how your body breaks down Juxtapid.
- Have moderate to severe liver problems or active liver disease, including abnormal liver function tests.

Other possible side effects of Juxtapid:

- The most common side effects of Juxtapid are stomach problems including diarrhea, nausea, vomiting, cramps/pain, indigestion, and/or gas. You may be able to reduce your chance of stomach problems by following an eating plan consisting of less than 20% of calories from fat.
- Juxtapid makes it harder for some fat-soluble nutrients, such as vitamin E and fatty acids, to get into your body. Take supplements that contain fatsoluble vitamins each day while you take Juxtapid. Ask your doctor, nurse, or dietitian how to take them.

Tell your doctor if you have any side effect that bothers you or that does not go away. Stop taking Juxtapid and tell your doctor if you have severe diarrhea, especially if you also have lightheadedness, decreased urine output, or tiredness. These are not all the possible side effects of Juxtapid. For more information, speak with your doctor or pharmacist.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Juxtapid may affect the way other medicines work, and other medicines may affect how Juxtapid works.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call 1-800-FDA-1088.

This is the most important information about Juxtapid. For more detailed information, Please see the Medication Guide and full Prescribing Information including Box Warning.







