

New Patient Start: Prescriber/Physician Order Form

PATIENT INFORMATION

Patient Name (Last, First) _____ Sex: Male Female Date of Birth ____/____/____ (mm/dd/yyyy)
 Address _____ City _____ State _____ ZIP _____
 Language: English Other _____
 Primary Contact Name _____ Relationship to Patient _____
 Primary Contact Email _____
 Primary Contact Phone _____ Primary Contact Cell _____

Please attach copies of patient insurance and prescription cards—front and back.

MEDICAL INFORMATION

Diagnosis: Alpha-mannosidosis (ICD-10-CM E77.1 Defects in glycoprotein degradation)
 Allergies: None Specify _____
Methods of Diagnosis (check all that apply):
 Enzyme Assay Genetic Testing Tissue Biopsy Other _____
 Prior treatment and dose: _____ Last date of prior treatment and dose: _____

Please attach copies of medical history/physical summary, urine oligosaccharides, acid α -mannosidase activity in leukocytes, current medications, genetic testing results, and allergies.

LAMZEDE® (VELMANASE ALFA-TYCV) PRESCRIPTION

Dosage — Available as Lamzede (velmanase alfa-tycv) 10 mg vial. Pharmacy to dispense quantity sufficient for a 28-day supply.
 Weight _____ lb or _____ kg Total Dose _____ mg* Route of Administration IV _____
 Frequency _____ Number of Refills _____

*The recommended dosage is 1 mg/kg (actual body weight) administered once weekly as an intravenous infusion.

Please list any additional treatment information, including follow-up evaluations: _____

PRESCRIBER/OFFICE INFORMATION

Prescriber's Name (Print) _____ Practice/Group Name _____
 Address _____ Suite _____
 City _____ State _____ ZIP _____
 Office Contact Person _____
 Office Phone _____ Office Fax _____
 License # _____ NPI # _____
 Preferred Medical Facility (Name, Phone) _____

List of Facilities Where Physician Has Privileges _____

By signing below, I certify that I am the prescribing provider mentioned above, that the therapy described above is medically necessary, and that all the medical necessity information is true, accurate, and complete. The patient's records contain supporting documentation that substantiates the utilization and medical necessity of the products marked above. I provide permission to use my personal information to facilitate this request and complete any regulatory or legal requirements associated with this request. I understand that the personal information provided herein may be shared with Chiesi, successors, and their agents and service providers as needed to support this request. I also attest that I have obtained the patient's authorization to release the above personal information and such other personal information as may be necessary for the Chiesi Total CareSM Program and/or their agents and service providers to facilitate this request and complete any regulatory or legal requirements associated with this request. If the patient is 18 years old or younger, I attest that I have obtained permission from the patient's legal guardian. If my patient is eligible for free product, I understand that receiving free product is not contingent on any purchase obligations. I also understand that no free product may be submitted for reimbursement to any payer, including Medicare and Medicaid; nor may I bill any payer for administration of such product. I understand that any falsification, omission, or concealment of material fact may result in criminal liability. For information on Chiesi USA, Inc.'s privacy practices, please view our Privacy Policy located at www.chiesiusa.com/privacy. Additionally, California residents may view Chiesi USA, Inc.'s California Notice at Collection at www.chiesiusa.com/privacy.

Dispense as written _____

 _____

Licensed Prescriber Signature (required – no stamps)

Printed Name _____ Date _____

ATTENTION: E-prescribe or use the official state prescription form where required by state law. No stamped signatures or signing on behalf of the prescriber.

Please continue on page 2. Please see Important Safety Information for Lamzede, including Boxed Warning for Severe Hypersensitivity Reactions, on page 2 and Full Prescribing Information.

New Patient Start: Prescriber/Physician Order Form (cont'd)

SITE OF SERVICE

Preferred Acquisitions Channel:

- Buy and Bill
(If site does not allow white bagging, send the form directly to EVERSANA®.)
- Specialty Pharmacy to Bill
Please send the prescription directly to:
EVERSANA Life Science Services
PH: 1-855-282-4883
FAX: 1-855-929-2828
- Other _____

Preferred Site of Infusion:

Site of Infusion _____
Contact Person _____

Indication

Lamzede® (velmanase alfa-tycv) is indicated for the treatment of non-central nervous system manifestations of alpha-mannosidosis in adult and pediatric patients.

Important Safety Information**WARNING: SEVERE HYPERSENSITIVITY REACTIONS****Hypersensitivity Reactions Including Anaphylaxis**

Patients treated with Lamzede have experienced hypersensitivity reactions, including anaphylaxis. Appropriate medical support measures, including cardiopulmonary resuscitation equipment, should be readily available during Lamzede administration. If a severe hypersensitivity reaction (e.g., anaphylaxis) occurs, discontinue Lamzede immediately and initiate appropriate medical treatment. In patients with severe hypersensitivity reaction, a desensitization procedure to Lamzede may be considered.

Considerations Due to Hypersensitivity Reactions and/or Infusion-Associated Reactions (IARs)

Prior to Lamzede administration, consider pretreating with antihistamines, antipyretics, and/or corticosteroids. Inform patients and caregivers of the signs and symptoms of hypersensitivity reactions and IARs and instruct them to seek medical care immediately if such symptoms occur.

- If a severe hypersensitivity reaction (including anaphylaxis) or severe IAR occurs, immediately discontinue Lamzede administration and initiate appropriate medical treatment.
- In the event of a mild to moderate hypersensitivity reaction or a mild to moderate IAR, consider temporarily holding the infusion for 15 to 30 minutes, slowing the infusion rate to 25% to 50% of the recommended rate, and initiating appropriate medical treatment.

Hypersensitivity Reactions Including Anaphylaxis

Anaphylaxis and severe hypersensitivity signs and symptoms included cyanosis, hypotension, emesis, urticaria, erythema, facial swelling, pyrexia, and tremor.

Infusion-Associated Reactions (IARs)

The most frequent symptoms of IARs that occurred in >10% of the population were pyrexia, chills, erythema, vomiting, cough, urticaria, rash, and conjunctivitis.

Females of Reproductive Potential

Advise females of reproductive potential to use effective contraception during treatment and for 14 days after the last dose if Lamzede is discontinued. For females of reproductive potential, verify that the patient is not pregnant prior to initiating treatment with Lamzede.

Embryo-Fetal Toxicity

Based on findings from animal reproduction studies, Lamzede may cause embryo-fetal harm when administered to a pregnant female.

Common Adverse Reactions

The most common adverse reactions (incidence >20%) are hypersensitivity reactions including anaphylaxis, nasopharyngitis, pyrexia, headache, and arthralgia.

Please see **Full Prescribing Information** for Lamzede.